

AD-A098 072

PITTSBURGH UNIV PA INST FOR STATISTICS AND APPLICATIONS F/6 12/1  
CHARACTERIZATIONS OF GENERALIZED MARKOV-POLYA AND GENERALIZED P--ETC(U)  
FEB 81 K G JANARDAN, B R RAO F89620-79-C-0161

UNCLASSIFIED TR-81-03

AFOSR-TR-81-0386

NL

1 of 1  
404  
098072

END  
DATE  
LSD  
5-81  
DTIC

AFOSR TR- 81-0386

LEVEL II

(2)

2  
AD A 09800

CHARACTERIZATIONS OF GENERALIZED MARKOV-POLYA AND  
GENERALIZED POLYA-EGGENBERGER DISTRIBUTIONS

K. G. Janardan\*

University of Pittsburgh  
and

B. Raja Rao

University of Pittsburgh

February 1981

Technical Report No. 81-03

Institute for Statistics and Applications  
Department of Mathematics and Statistics  
University of Pittsburgh  
Pittsburgh, PA 15260

This document is best quality print quality.  
The copy furnished to DDC contained a  
significant number of pages which do not  
exceed legibly.

\*On Sabbatical leave from Sangamon State University, Springfield, IL 62708

DTIC  
ELECTED  
APR 23 1981  
S F D

FILE COPY  
DTIC

Part of this work is sponsored by the Air Force Office of Scientific Research under Contract [REDACTED]. Reproduction in whole or in part is permitted for any purpose of the United States Government.

F49620-79-C-0161

## **DISCLAIMER NOTICE**

**THIS DOCUMENT IS BEST QUALITY  
PRACTICABLE. THE COPY FURNISHED  
TO DTIC CONTAINED A SIGNIFICANT  
NUMBER OF PAGES WHICH DO NOT  
REPRODUCE LEGIBLY.**

## REPORT DOCUMENTATION PAGE

UNCLASSIFIED

READ INSTRUCTIONS  
BEFORE COMPLETING FORM  
DO NOT OVERWRITE OR ERASE1. REPORT NUMBER  
**AFOSR-TR-81-0386**2. GOVERNMENT ACQUISITION NO.  
**4D-AC98072**3. TITLE (Include Subtitle)  
**Characterizations of Generalized Markov-Polya and  
Generalized Polya-Eggenberger Distributions**

4. NAME OF REPORT SOURCE

**INTERIM**5. PREVIOUS EDITION NUMBER  
**U.P. Study Report 61-11**

6. DISTRIBUTION STATEMENT

**F49620-79-C-0161**

7. AUTHOR(S)

**K. G. Janardan & P. Raja Rao**

8. PERIODICITY/REPORT DATE

9. COLLECTING/STUDY NUMBER  
**2304/95**University of Pittsburgh  
Dept. of Math. & Statistics  
Pittsburgh, PA 15260**61102F**

10. CONTROLLING OFFICE NAME AND ADDRESS

**AFOSR/NM  
Bolling AFB, DC 20332**

11. APPROVING AUTHORITY

February 1981

12. APPROVAL NUMBER

13

13. APPROVAL DATE

14. APPROVAL COMMENTS

15. APPROVAL SIGNATURE

Approved for public release;  
distribution unlimited.

16. DESTROY DATE OR DURATION OF THIS REPORT

17. SECURITY CLASSIFICATION

18. SUBJECT TERMS (List subjects which best describe the content of the report)

Generalized Chapman-Dorfman's distributions, generalized Markov-Polya distributions, characterizations, conditional distributions, characterizations, Markov-Polya distributions.

19. KEY WORDS (List words or terms which describe the content of the report)

A generalization of the Chapman-Dorfman's distributions is introduced. The properties of the distributions are studied. The distributions are shown to be partially exchangeable and hence the moments and cumulants of the distributions are obtained. The characterization of the generalized Chapman-Dorfman's distributions is obtained. The distributions are shown in the context of the three main conditions. Other topics such as tail ordering theorems are also proved concerning the probability distributions.

DD FORM 1473

1 APR 1973  
GSA GEN. REG. NO. 27, 1473-100-02

CHARACTERIZATIONS OF GENERALIZED MARKOV-POLYA AND  
GENERALIZED POLYA-EGGENBERGER DISTRIBUTIONS

K. G. Janardan\* and B. Raja Rao  
Department of Mathematics & Statistics  
University of Pittsburgh  
Pittsburgh, PA 15260

ABSTRACT.

A discrete model is considered where the original observation is subjected to partial destruction according to the generalized Markov-Polya damage model. A characterization of the generalized Polya-Eggenberger distribution is given in the context of the Rao-Rubin condition. Several other characterization theorems are also proved concerning these probability distributions

Key Words & Phrases: Generalized Markov-Polya distribution; generalized Polya-Eggenberger distribution; damaged model; conditional distribution characterizations.

Accession For	
NTIS GRA&I	X
DTIC TAB	
Unannounce	
Justificat	
By _____	
Distributi	
Availability Codes	
Avail and/or	
Dist	Special
A C M	

\* On Sabbatical leave from Sangamon State University  
Springfield, IL 62703

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF THIS INITIAL TO DDC  
This technical document has been reviewed and is  
approved for public release by AFSC AFM 1AFR 100-12 (7b).  
Distribution is unlimited.  
A. D. [Signature]  
Technical Information Officer

## 1. INTRODUCTION

Urn models are readily adapted to the development of probability distributions used in the analysis of complex problems in real life situations. In many natural phenomena involving individuals or living organisms, the probability of success seems to increase or decrease with the number of successes or failures. Thus, with the aid of urn models, research workers have developed a number of discrete probability distributions (see Johnson and Kotz, 1977; chp. 4) when the probability of success of an event is a linear function of the number of successes. Among the principal researchers who have used urn models for developing discrete probability models for contagious events, Markov (1917) and Polya and Eggenberger (1923) are pioneers in the field.

Following Hald's (1960) approach, Janardan (1973), and Janardan and Schaeffer (1977) have recently considered a new three urn model with predetermined strategy and have derived the generalized Markov-Polya distribution (GMPD) as a model (1.1) for voting in small groups where contagion is present within each group and the group leader devises some new strategies for bringing success to his candidate:

$$P(X=x) = \binom{n}{x} \frac{a^b b(a+b+nt)}{(a+b)(a+xt)(b+(n-x)t)(a+bt+nt)} \frac{(x,c)}{(b+(n-x)t)} \frac{(n-x,c)}{(a+bt+nt)} \quad (1.1)$$

where  $a>0$ ,  $b>0$ ,  $0 \leq t \leq 1$ ,  $c \neq 0$ , and  $x = 0, 1, 2, \dots, n$ . If  $t = 0$  this reduces to Markov-Polya distribution (see equation (4.1) of Johnson-Kotz, 1977, p. 177). There is a connection the Polya-Eggenberger distribution.

The GMPD (1.1) has several interesting properties and a large number of applications (see Janardan and Schaeffer, 1977 and Janardan, 1978). Under certain limiting conditions, the probability distribution (1.1) gives the limiting form:

$$P(X=x) = \frac{h(0)xt^x}{(h!x!)x!c} \cdot p^x (1-p)^{h(x-1)/c} \quad (1.2)$$

where  $x=0,1,2,\dots$ ,  $b>0$ ,  $0<\beta<1$ ,  $0\leq t\leq 1$  and  $c\neq 0$ .

This distribution was named "The generalized Polya-Eggenberger" distribution (GPE) by Janardan (1973) since (1.2) reduces to the Polya-Eggenberger distribution when  $t=0$  (see Patil and Joshi, 1968 p. 20).

Situations often arise where the original observation produced by nature undergoes a destructive process and what is recorded is only the damaged portion of the actual (original) observation. This problem was first brought to light by Sie (1963) when he considered the resultant models after the observations, produced by some probability model, were ruined by other probability models. Subsequently, Rao and Rubin (1964) proved that if the observation generated by nature (denoted as r.v. $X$ ) is reduced to  $Y$  by a binomial destructive process and if  $Y$  satisfies the condition:

$$P(Y=k) = P(Y=k/\text{no damage}) = P(Y=k/\text{partial damage}), \quad (1.3)$$

then the original r.v. $X$  must have the Poisson distribution. In the literature, this result is known as Rao-Rubin characterization of the Poisson distribution and the condition (1.3) is called RR-condition.

In this paper, we consider the generalized Markov-Polya distribution as a damage model subject to RR condition and characterize the generalized Polya-Eggenberger distribution (GPE) as a model of contagion for the production of observations in nature. In addition to this result which is given in the next section, we prove several other theorems which characterize the generalized Markov-Polya and generalized Polya-Eggenberger distributions.

## 2. NOTATION AND IDENTITIES

To begin with, we shall discuss the notation and two identities required in this paper. The notation  $\pi_{(x,c)}$  used in the definitions of probability distributions (1.1) and (1.2), and in sequel stands for the ascending factorials:

$$\begin{aligned}
 m(x, c) &= m(m+c)(m+2c)\dots(m+(x-1)c), \\
 m(0, 0) &= m^0, \quad m(0, c) = 1, \\
 m(x, -1) &= m(x) = m(m-1)\dots(m-x+1), \\
 m(x, +1) &= m[x] = m(m+1)\dots(m+x-1).
 \end{aligned} \tag{2.1}$$

From Janardan (1973), we record the following two identities:

$$\sum_{k=0}^n J_k(a, t, c) J_{n-k}(b, t, c) = J_n(a+b, t, c) \tag{2.2}$$

$$\text{where } J_m(a, t, c) = a(a+mt)^{(m, c)} / (a+mt)m! \tag{2.3}$$

$$\text{and } J_0(a, b, c) = 1. \tag{2.4}$$

$$\sum_{k=0}^{\infty} J_k(a, t, c) V^k = U^{-A/c}, \quad \text{with } V = (1-U)U^t/c.$$

These identities can be derived by using Lagrange's expansion.

With this notation, the probability distributions (1.1) and (1.2) can be written respectively as

$$P(X=x) = J_x(a, t, c) J_{n-x}(b, t, c) / J_n(a+b, t, c), \tag{2.5}$$

$$P(X=x|Y=y) = J_x(a, t, c) (y/c)^x (1-y)^{(b+xt)/c} \tag{2.6}$$

### 3. CHARACTERIZATIONS BASED ON CONDITIONAL DISTRIBUTIONS

We now prove the following theorem:

**THEOREM 1:** If  $X$  and  $Y$  are two independent random variables having the generalized Polya-Eggenberger distributions with parameters  $(a, t, c, \beta)$  and  $(b, t, c, \beta)$  respectively, then the conditional distribution of  $X$  given  $X+Y=n$  is a generalized Markov-Polya distribution as given in (1.1).

**PROOF:** By definition of conditional probability,

$$P(X=x|X+Y=n) = P(X=x, Y=n-x) / P(X+Y=n)$$

$$\begin{aligned}
&= \frac{\sum_{x=0}^n [J_x(a, t, c) (\beta/c)^x (1-\beta)^{(a+xt)/c}] [J_{n-x}(b, t, c) (\beta/c)^{n-x} (1-\beta)^{(b+(n-x)t)/c}]}{\sum_{x=0}^n [J_x(a, t, c) (\beta/c)^x (1-\beta)^{(a+xt)/c}] [J_{n-x}(b, t, c) (\beta/c)^{n-x} (1-\beta)^{(b+(n-x)t)/c}]} \\
&= \frac{J_x(a, t, c) J_{n-x}(b, t, c)}{\sum_{x=0}^n J_x(a, t, c) J_{n-x}(b, t, c)}
\end{aligned}$$

By identity (2.2), the denominator equals  $J_n(a+b, t, c)$ , which proves the theorem. The following theorem gives the converse of the above.

THEOREM 2: Let X and Y be two independent discrete r.v's such that the conditional distribution of  $X=x$  given  $X+Y=n$  is the generalized Markov-Polya distribution given by (1.1) or (2.3) then each X and Y has a generalized Polya-Iggenberger distribution as in (1.2).

PROOF: By hypothesis of the theorem,  $P(X=x|X+Y=n)$  is given by

$J_x(a, t, c) J_{n-x}(b, t, c) J_n(a+b, t, c)$  which is of the form

$\alpha(x)\beta(n-x)/\gamma(n)$  with  $\alpha(x) = J_x(a, t, c)$ ,  $\beta(n-x) = J_{n-x}(b, t, c)$  and  $\gamma(n) = J_n(a+b, t, c)$ .

Applying theorem 1 of Janardan (1975), we get

$f(x) = p\alpha(x)e^{rx}$  and  $g(y) = q\beta(y)e^{ry}$ , where p, q and r are some positive constants. Setting  $e^r = \beta(1-\beta)^{1/c}$ , the functions f(.) and g(.) can be written as

$$f(x) = p J_x(a, t, c) \beta^x (1-\beta)^{ax/c}, \text{ and}$$

$$g(y) = q J_y(b, t, c) \beta^y (1-\beta)^{by/c}.$$

Since  $1 = \sum_{x=0}^{\infty} f(x) = \sum_{y=0}^{\infty} g(y)$ , applying the identity (2.4), we get

$p = (1-\beta)^{-a/c}$  and  $q = (1-\beta)^{-b/c}$  completing the proof of the theorem.

THEOREM 3 : If a non-negative integer random variable  $N$  is sub-divided into two components  $N_A$  and  $N_B$  such that the conditional distribution  $P(N_A = x, N_B = n-x|N=n)$  is the generalized Markov-Polya distribution (2.5) then the r.v's  $N_A$  and  $N_B$  are independent if, and only if,  $N$  has a generalized Polya-Eggenberger distribution.

PROOF: The joint probability of  $N_A$  and  $N_B$  becomes

$$P[N_A=x, N_B=n-x] = \frac{J_x(a, t, c) J_{n-x}(b, t, c)}{J_n(a+b, t, c)} \cdot P(N=n). \quad (3.1)$$

If  $N$  has a generalized Polya-Eggenberger distribution, then with  $h = a+b$ , its probability function is

$$P(N=n) = J_n(a+b, t, c) (\beta/c)^n (1-\beta)^{(a+b+nt)/c}, \quad (3.2)$$

$$a>0, b>0, 0 \leq t \leq 1, 0 < \beta < 1, c \neq 0.$$

Inserting the value of  $P(N=n)$ , we can easily write (3.1) as

$$\begin{aligned} P[N_A=x, N_B=n-x] &= \\ J_x(a, t, c) (\beta/c)^x (1-\beta)^{(a+xt)/c} J_{n-x}(b, t, c) (\beta/c)^{n-x} (1-\beta)^{(b+(n-x)t)/c} \\ &= P(N_A=x) P(N_B=n-x). \end{aligned}$$

Conversely, if  $N_A$  and  $N_B$  are independent r.v's such that the conditional distribution of  $N_A$  and  $N_B$  given  $N_A + N_B = n$  is the GMFD, then  $N_A$  and  $N_B$  have the generalized Polya-Eggenberger distributions. This follows from theorem 2.

THEOREM 4: If  $X$  and  $Y$  are two independent r.v's defined on non-negative integers such that  $P(X=x) = f(x) > 0, f(0) = 1$  and

$$x \geq 0$$

$$P(Y=y) = g(y) > 0,$$

$$\sum_{y=0}^{\infty} g(y) = 1 \text{ and further if for } \frac{a_n}{b_n} = A,$$

$$P(X=k/X+Y=n) = \frac{\binom{n}{k} a_n^k b_n^{n-k} (a+nt)^{(k,c)} (b+nt)^{(n-k,c)}}{(a+nt)^{(n,c)} (b+nt)^{(n,c)}} \quad (3.3)$$

for  $k = 0, 1, 2, \dots, n$ .

$= 0$  for  $k > n$

Then (i)  $a_n$  is independent of  $n$  and equals a constant 'a' for all values of  $n$ , and

(ii)  $X$  and  $Y$  must have generalized Polya-Eggenberger distributions with parameters  $(a, t, c, s)$  and  $(b, t, c, r)$  respectively.

PROOF: Since  $X$  and  $Y$  are independent r.v's we have

$$P(X=k/X+Y=n) = f(k)g(n-k)/ \sum_{k=0}^n f(k)g(n-k) . \quad (3.4)$$

Using (3.3) and (3.4) we can derive the functional relation (3.5) for all values of  $0 \leq k \leq n$ , and  $n \geq 1$ :

$$\frac{f(k)g(n-k)}{f(k-1)g(n-k+1)} = \frac{n-k+1}{k} \frac{(a_n+kt)^{(k,c)} (b_n+nt)^{(n-k,c)}}{(a_n+(k-1)t)^{(k-1,c)} (b_n+(n-k+1)t)^{(n-k,t)}} \frac{(a_n+(k-1)t)^{(k-1,t)}}{(a_n+kt)^{(k,t)} (b_n+nt)^{(n-k,t)}} \quad (3.5)$$

Replacing  $k$  by  $k+1$  and  $n$  by  $n+1$  in (3.5) and dividing the resulting expression by (3.5) we get  $f(k+1)f(k-1)/f^2(k)$  on the left side and a very complex untidy expression on the right side. Since the left side is independent of  $n$ , the right side must also be independent of  $n$ . Thus,  $a_n$  and  $b_n$  must be the same for all  $n$ , and hence ignoring the subscripts on  $a$ 's and  $b$ 's we will

have  $\frac{f(k+1)f(k-1)}{f^2(k)} =$

$$\frac{k}{k+1} \frac{(a+(k-1)t)^{(k-1,c)} (a+(k+1)t)^{(k+1,c)}}{(a+(k-1)t)(a+(k+1)t)[(a+k)t]} \frac{(a+k)^2}{(k,c)} , \quad (3.6)$$

which by continued substitution for  $k=1, 2, \dots, (n-1)$ , and multipli-

lication together yields

$$\frac{f(n)f(0)}{f(n-1)f(1)} = \frac{1}{n!} \frac{(a+nt)^{(n,c)} (a+(n-1)t)^{(n-1,c)}}{a(a+nt)(a+(n-1)t)^{(n-1,c)}}. \quad (3.7)$$

Setting  $B = f(1)/f(0)a$ , the recurrence relation (3.7) becomes

$$f(n) = \frac{B}{n!} \frac{(a+nt)^{(n,c)} (a+(n-1)t)^{(n-1,c)}}{(a+nt)(a+(n-1)t)^{(n-1,c)}} f(n-1), \quad (3.8)$$

which is true for all integral  $n$ . Thus,

$$f(n) = B^n a(a+nt)^{(n,c)} f(0) / (a+nt) n!. \quad (3.9)$$

Since  $\sum_{n=0}^{\infty} f(n) = 1$ , the series (3.9) must converge to unity. Let

the unknown positive quantity  $B$  be equal to  $(\beta/c)(1-\beta)^{t/c}$ ,  $0 < \beta < 1, t > 0$ , and  $c \neq 0$ . Thus,

$$1 = \sum_{n=0}^{\infty} \frac{a(a+nt)^{(n,c)} n!}{(a+nt)n!} \frac{(\beta/c)}{(1-\beta)^{nt/c}} f(0).$$

By applying identity (2.4), we can easily see that  $f(0) =$

$$(1-\beta)^{-a/c} \text{ and } f(x) = \int_x^\infty (a, t, c) (\beta/c)^x (1-\beta)^{(x+tx)/c},$$

which proves that the r.v.  $X$  is distributed as the GPED with parameters  $(a, t, c, \beta)$ . By putting  $x=0$  in (3.5), one can easily see that

$$\frac{g(n)}{g(n-1)} = \frac{B(b+nt)^{(n,c)} (b+(n-1)t)^{(n-1,c)}}{n! (b+nt)(b+(n-1)t)^{(n-1,c)}}.$$

Hence  $g(n) = B^n b(b+nt)^{(n,c)} g(0) / (b+nt)n!$

and the fact  $\sum_{n=0}^{\infty} g(n) = 1$  will similarly give  $g(0) = (1-\beta)^{-b/c}$  for

$B = (\beta/c)(1-\beta)^{t/c}$ . Thus, the r.v.  $Y$  must be the GPED with parameters  $(b, t, c, \beta)$ .

REMARK: It was shown in theorem 3 , that if  $X$  and  $Y$  are independent generalized Polya-Eggenberger random variables, then the conditional distribution of  $X$  given  $X+Y$  is generalized Markov-

**Polya distribution.** The above theorem, which was motivated by theorem 1 of Chatterji (1963), shows that a weak form of this property characterizes the r.v. X.

#### 4. CHARACTERIZATION THEOREMS BASED ON THE IN CONDITION

Let  $(X, Y)$  be a random vector of non-negative integer-valued components such that

$$P(X=n, Y=k) = f_n S(k/n) \quad (4.1)$$

where  $\{f_n : n=0,1,2,\dots\}$  and  $\{S(k/n) : k=0,1,2,\dots,n\}$  for each  $n \geq 0$  are discrete probability distributions. That is, the marginal distribution of X is  $\{f_n\}$  and for each  $n \geq 0$  with  $f_n > 0$ , the conditional distribution of Y given  $X=n$  is  $\{S(k/n) : k=0,1,2,\dots,n\}$ . Further,

$$P(Y=k/\text{no damage}) = f_k S(k/n) / \sum_{j=0}^n f_j S(j/n) \quad (4.2)$$

$$P(Y=k/\text{damaged}) = \sum_{n=k+1}^{\infty} f_n S(k/n) / \sum_{n=k+1}^{\infty} f_n S(n/n). \quad (4.3)$$

**THEOREM 5:** If a r.v. X defined on non-negative integers is distributed in nature as a GPFD (1.1) with parameters  $(a, t, c, \beta)$  and if it is damaged and reduced to k by the GMD (1.1) and further, if Y is the resulting random variable, then (i) the Condition (1.3) is satisfied, and (ii) has a GMD with parameters  $(a, t, c, \beta)$ .

PROOF:  $P(Y=k) = \sum_{n=k}^{\infty} f_n S(k/n)$  because  $P(Y=k) =$   

$$\sum_{n=k}^{\infty} [J_n(a+b, t, c)(\beta/c)^n (1-\beta)^{(a+b+t)/c}] J_k(a, t, c) J_{n-k}(b, t, c) J_n(a+b, t, c)]$$
  
 $= J_k(a, t, c)(\beta/c)^k (1-\beta)^{(a+b+t)/c} \sum_{n=k}^{\infty} J_{n-k}(b, t, c)(\beta/c)^{n-k} (1-\beta)^{(a+b+(n-k)t)/c}]$   
 $= J_k(a, t, c)(\beta/c)^k (1-\beta)^{(a+b+t)/c}$  since the sum of the terms in the square brackets is one.

From equation (4.2),

$$P(Y=k/\text{no damage}) = J_k(a, t, c)(t/c)^k (1-t/c)^{(a+b)/c} = P(Y=k).$$

From equation (4.3),

$$\begin{aligned} P(Y=k/\text{damaged}) &= \frac{\sum_{n=k}^{\infty} J_n(a, t, c)(t/c)^k (1-t/c)^{(a+b)/c}}{\sum_{k=0}^{\infty} J_k(a, t, c)(t/c)^k (1-t/c)^{(a+b)/c}} \\ &= P(Y=k). \end{aligned}$$

THEOREM 6: Let  $X$  be a non-negative integer-valued r.v. and let the probability that an observation  $n$  of  $X$  is reduced to  $k$  during a destructive process be given by the GMPP:

$$S(k/n) = J_k(a, t, c)J_{n-k}(b, t, c)/J_n(1, t, c),$$

for  $0 < a < 1$ ,  $a+b=1$ ,  $0 \leq t \leq 1$ ,  $c > 0$ , and  $k=0, 1, 2, \dots, n$ . If the resulting r.v.  $Y$  is such that it satisfies the RR-Condition (1.3), then  $X$  has a GPED with parameters  $(1, t, c, f)$ .

PROOF: The RR-Condition is equivalent to

$$\sum_{n=k}^{\infty} f_n S(k/n) = f_k S(k/k) / \sum_{j=0}^{\infty} f_j S(j/j),$$

where  $f_k = P(X=k)$ . This gives

$$\sum_{n=k}^{\infty} f_n \frac{J_k(a, t, c)J_{n-k}(b, t, c)}{J_n(1, t, c)} = \frac{f_k J_k(a, t, c)/J_k(1, t, c)}{\sum_{j=0}^{\infty} f_j J_j(a, t, c)/J_j(1, t, c)}. \quad (4.4)$$

setting  $n=k+s$  and cancelling  $J_k(a, t, c)$  on both sides we get

$$\sum_{s=0}^{\infty} f_{k+s} \frac{J_s(b, t, c)}{J_{k+s}(1, t, c)} = \frac{f_k/J_k(1, t, c)}{\sum_{j=0}^{\infty} f_j J_j(a, t, c)/J_j(1, t, c)}. \quad (4.5)$$

$$\text{Define } f_k = F(k) J_k(1, t, c) v^k, \quad (4.6)$$

for all integral values of  $k$ , where  $v$  is some arbitrary quantity to be set later. Substituting (4.6) in (4.5) we get

$$\sum_{s=0}^{\infty} F(k+s) J_s(b, t, c) v^{k+s} = F(k) v^k / \sum_{j=0}^{\infty} F(j) J_j(a, t, c) v^j \quad (4.7)$$

$$\text{Let } G(az, t) = \sum_{k=0}^{\infty} F(k) J_k(az, t, c) v^k,$$

where  $-\infty < z < \infty$  so that  $G(0, t) = F(0)$  and  $G(1, t) = 1$ .

Multiplying both sides of (4.7) by  $J_k(az, t, c)$  and summing over  $k$  from 0 to  $\infty$  (4.8) becomes

$$\sum_{n=0}^{\infty} F(n) v^n \left\{ \sum_{k=0}^n J_k(az, t, c) J_{n-k}(b, t, c) \right\} = G(az, t) / G(a, t). \quad (4.9)$$

By identity (2.2) the inner sum on the left side of (4.9) is equivalent to  $J_n(az+b, t, c)$ , and hence (4.9) gives the bivariate functional equation:

$$G(az+b, t) G(a, t) = G(az, t). \quad (4.10)$$

Clearly  $G(a+b, t) = G(1, t) = 1$ . Setting  $x=a(z-1)$  and  $b=1-a$  (4.10) gives

$$G(x+1, t) G(a, t) = G(x+a, t) \quad (4.11)$$

Setting  $\phi(x) = G(x+1, t)$ , and  $a=1-x$  (4.11) reduces to the Cauchy functional equation,  $\phi(x)\phi(y) = \phi(x+y)$ , whose non-trivial solution is given by  $\phi(x) = e^{\lambda x}$ . Thus,  $G(x, t) = e^{\lambda(x-1)}$  which is the probability generating function of the Poisson distribution.

Now replacing  $x$  by  $az$ , assigning a value  $(1-e^{-\lambda c})e^{-\lambda t}/c$  to  $v$ , using the definition of  $G(az, t)$ , we get

$$e^{\lambda az} = \sum_{k=0}^{\infty} F(k) e^{\lambda} J_k(az, t, c) v^k \quad (4.12)$$

To determine the value of  $F(k)$ , consider the identity (2.4) with  $A = az$  and  $w = e^{-\lambda c}$  which gives

$$e^{\lambda az} = \sum_{k=0}^{\infty} J_k(az, t, c) v^k . \quad (4.13)$$

Subtracting (4.13) from (4.12),

$$0 = \sum_{k=0}^{\infty} [F(k)e^{\lambda} - 1] J_k(az, t, c) v^k . \quad (4.14)$$

Since (4.14) is true for all values of  $\lambda$ , it is obvious that  $F(k)=e^{-\lambda}$  for all  $k$ . Hence, by definition (4.6), we get

$$\begin{aligned} f(k) &= F(k) J_k(1, t, c) v^k \\ &= e^{-\lambda} J_k(1, t, c) [e^{-\lambda t} (1-e^{-\lambda c})/c]^k \\ &= J_k(1, t, c) (\beta/c)^k (1-\beta)^{(1+k)t}/c \\ &\quad \text{with } \beta=1-e^{-\lambda c} . \end{aligned}$$

That is,  $X$  has a GPED with parameter  $(1, t, c, \beta)$ .

#### ACKNOWLEDGEMENTS

The first author thanks professor P.R.Krishniah for all the facilities provided to him while the author was spending his sabbatical leave at the University of Pittsburgh. Miss Beth Ford typed the manuscript.

BIBLIOGRAPHY

- (1). Chatterji, S. D. (1963). Some elementary characterizations of the Poisson distribution. Amer. Math. Monthly 70, 953-64.
- (2). Hald, A. (1960). The Compound hypergeometric distribution and a system of single sampling inspection plans based on prior distributions and costs. Technometrics 2, 275-340.
- (3). Janardan, K. G. (1973). A four urn model with predetermined strategy. Technical Report, Sangamon State University, Springfield, IL
- (4). Janardan, K. G. (1975). Characterizations of certain discrete distributions. Statistical Distributions in Scientific Work 3, (Eds. Patil et.al) D. Reidel Publishing Co. Holland. pp. 359-64.
- (5). Janardan, K. G. (1978). On a generalized Markov-Polya distribution. Gujarat Statistical Review 5 (1), 16-32.
- (6). Janardan, K. G. & Schaeffer, D. J. (1977). A generalization of Markov-Polya distribution: Its extensions and applications. Biom. J 19 (2), 87-106.
- (7). Johnson, N. L. & Kotz, S. (1977). Urn models and their applications. John Wiley & Sons Inc., New York and Canada.
- (8). Markov (1917). On limiting formulas for computing probabilities (O. Nekotovih Predelnih Formulah Icislenia Veroiatnostei-Russian). Izvestia Imperatorsky Academy Nauk, Seria VI, II (3), 177-36.
- (9). Patil, G. P. & Joshi, S. W. (1968). A Dictionary and Bibliography of Discrete Distributions, Oliver & Boyd, Edinburgh.
- (10). Polya, G. & Eggenberger, F. (1923). Über die Statistik verketteter vorgänge. Zeitschrift fur Angewandte Mathematik und Mechanik 3, 279-89.
- (11). Rao, C. R. (1963). On discrete distributions arising out of methods of ascertainment. Int. Symp. Discrete Distributions, Montreal.
- (12). Rao, C. R. & Rubin, H. (1964). On a characterization of the Poisson distribution. Sankhya A 26, 295-98.

